



Abamec LV Pour-On for Cattle



Vetsense Abamec LV Pour-On for Cattle is ideal for the treatment and control of Abamectin sensitive internal and external parasites of dairy and beef cattle.

Contains Abamectin (10 g/L), a member of the macrocyclic lactones.

- ✓ Low-Volume formula (double strength)
- ✓ Contains Abamectin
- ✓ Easy application
- ✓ Zero-day milk withholding period
- ✓ Controls both internal and external parasites

Abamec LV Pour-On

ACTIVE CONSTITUENTS	10 g/L ABAMECTIN
SPECIES	Cattle - Beef & Dairy
DOSE RATE	1 mL / 20 kg
ADMINISTRATION	Pour-On
PACK SIZES	1L, 2.5L, 5L & 10L
APVMA NO	95453/145718
WITHHOLDING PERIOD	
MEAT	35 days
MILK	Nil
EXPORT SLAUGHTER INTERVAL	An ESI has not been established for this product.



Contact
Vetsense:



PRODUCT INFORMATION

Abamec LV Pour-On for Cattle

Active Constituents: 10 g/L ABAMECTIN



Vetsense Abamec LV Pour-On for Cattle contains Abamectin (10 g/L), a member of the macrocyclic lactones.

Abamec LV is effective against abamectin sensitive strains of internal and external parasites when applied to the skin along the backline of cattle. Resistance may develop to any chemical.

INDICATIONS

For the treatment and control of adult and immature stages of:

✓ GASTRO-INTESTINAL ROUNDWORMS

- Barber's pole worm (*Haemonchus placei*)
- Small brown stomach worm, including inhibited stages (*Ostertagia ostertagi*)
- Small intestinal worm (*Cooperia punctata*)
- Intestinal nematode (*Cooperia oncophora*)
- Stomach hair worm (*Trichostrongylus axei*)
- Hookworm (*Bunostomum phlebotomum*)
- Nodule worm (*Oesophagostomum radiatum*)
- Whipworm (*Trichuris spp.*)
- Thin necked intestinal worm (*Nematodirus spathiger*)
- Intestinal threadworm (*Strongyloides papillosus*) and Large lungworm (*Dictyocaulus viviparus*)

✓ SUCKING AND BITING LICE

- Long nosed sucking louse (*Linognathus vituli*)
- Short nosed sucking louse (*Haematopinus eurysternus*)
- Biting louse (*Bovicola bovis*)
- Little sucking louse (*Solenopotes capillatus*)

✓ CATTLE TICK (*Boophilus microplus*)

- Aids in control (Not recommended in strategic dipping programs for cattle tick control).

SUSTAINED ACTIVITY

At the recommended dose rate, Vetsense Abamec LV Pour-On for Cattle effectively controls infection with *Ostertagia spp.* (Small brown stomach worm) acquired up to 14 days after treatment, with *Dictyocaulus viviparus* (Lungworm) acquired up to 28 days after treatment.

CONTRAINDICATIONS

This product is contraindicated for use in calves under 16 weeks of age or less than 50 kg body weight.

This product should not be used in dogs, horses or any other pets.

This product is not recommended for use in goats as safety and efficacy have not been evaluated.

DOSAGE AND ADMINISTRATION

Apply to the skin along the backline of the cattle at the dose rate of 1 mL of Vetsense Abamec LV Pour-On for Cattle per 20 kg bodyweight.

Cattle heavier than 650 kg should be dosed at 1 mL per 20 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not underdose. Where there is a large variation in size within the group, draft into 2 or more lines based on body weight, to avoid excessive overdosing.

GENERAL DIRECTIONS

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 35 days before slaughter for human consumption.

MILK: Zero (0) Days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not been established for this product.

NOTE: Observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Vetsense on 0245779911 before using this product.

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

ADDITIONAL USER SAFETY

Additional information is listed in the Safety Data Sheet (SDS).

ENVIRONMENTAL PROTECTION

Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical and used containers.

DISPOSAL

For containers equal to 1L:

Dispose of container by wrapping with paper and putting in garbage.

For large containers greater than 1L:

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature) in the closed, original container, in a dry, well-ventilated area, and protect from sunlight.

APVMA No.: 95453/145718

LIVWEIGHT	DOSE VOLUME	1L PACK TREATS	2.5L PACK TREATS	5L PACK TREATS	10L PACK TREATS
51-100 kg	5 mL	200	500	1000	2000
101-150 kg	7.5 mL	133	333	666	1333
151-200 kg	10 mL	100	250	500	1000
201-250 kg	12.5 mL	80	200	400	800
251-300 kg	15 mL	66	166	333	666
301-350 kg	17.5 mL	57	142	285	571
351-400 kg	20 mL	50	125	250	500
401-450 kg	22.5 mL	44	111	222	444
451-500 kg	25 mL	40	100	200	400
501-550 kg	27.5 mL	36	90	181	363
551-600 kg	30 mL	33	83	166	333
601-650 kg	32.5 mL	30	76	153	307

1L-Pack Application Guide

STEP 1

Turn bottle upside down to fill the dosing chamber.




STEP 2

Once chamber is filled, stand bottle upright to level the dosage.



STEP 3

Tilt bottle to adjust to the correct dosage.



STEP 4

Once you have the correct dosage, remove the cap.



STEP 5

Pour out the liquid.



STEP 6

Apply cap to reseal the bottle.

